SOME JUNE-JULY BIRD RECORDS FOR THE PRAIRIE PROVINCES

. BERNARD GOLLOP, Canadian Wildlife Service, 115 Perimeter Rd., Saskatoon, askatchewan. S7N 0X4

This article was originally written for merican Birds, published six times a year y the National Audubon Society, 950 Third we., New York, N.Y. 10022. Four issues over reports of outstanding bird events moughout North America in spring, summer, all and winter. Hundreds of cooperators ontribute records to compilers of data for 26 egions, including the Prairie Provinces egion. Of the other two issues, one is evoted to breeding and wintering bird cenuses and the other to Christmas Bird Counts - 1357 in 1981. The six issues, which also ontain articles on birds, exceed 1000 ages. Subscription is \$18 (US).

For Saskatchewan, reports are usually ceived from natural history societies in loose Jaw, Regina and Saskatoon and from irdwatchers living in Biggar, Broadview, arnduff, Dilke, Eastend, Foam Lake, Froner, Good Spirit Lake, Grenfell, Indian Head, enaston, Kenosee Lake, Kindersley, La oche, Oxbow, Raymore, Spring Valley, Val larie, Waskesiu Lake, Webb, White Bear nd Wolseley. A separate report is compiled om these data. (See Jim Wedgwood's sumnary of spring migration in this issue.)

Anyone wishing to become part of this ctwork should contact Bernie Gollop. otices and forms will then be sent out. The inter period covers Dec. 1 through Feb. 28. pecies are discussed in A.O.U. order.

Mean temperatures in June 1982 anged from 2.5°C above normal in entral Alberta to 3°C below average at hurchill and Winnipeg. July mperatures were near normal across e region. Precipitation was scanty verywhere in June. Edmonton reported s second-driest June on record and any locations in Saskatchewan were e driest in more than 20 years. ochrane, Alta., had 4 in. of snow 5-6 une. July precipitation was almost normal in Saskatchewan and up to twice normal west of Edmonton and north of Winnipeg. Lethbridge, Alta., had 63 mm rain in less than 1 hour and Emerson, Man., had 81 mm in 75 minutes.

July ponds in s. Alberta totalled 503,000 (54% above last year), 964,000 in s. Saskatchewan (165% more than last July) and 195,000 in s. Manitoba (7% better than a year ago).

On 29 May from Saskatoon w. and s. in Saskatchewan (and into s. Alberta), there was a snowstorm that dumped 10 cm (4 in.) of snow around Saskatoon with average winds of 38 km/hour and a temperature range of 0°-2.6°C. In the Cypress Hills area drifts reached 6 ft. and thousands of livestock were lost. Near Saskatoon, many swallows rested on the snow while others were apparently feeding over open water. Much of the snow disappeared within 3 days but the effect on birds, particularly insectivorous species, must have been devastating. However, no one reported numbers of dead birds. Barn Swallows, Least Flycatchers, House Wrens and Clay-colored Sparrows decreased significantly after the storm around Swift Current (CH). See also Tree Swallow and Mountain Bluebird accounts below.

LOONS THROUGH WATERFOWL — Thirty Com. Loons had congregated on Waskesiu L., Prince Albert N.P., by 25 June (S. Ross). A Red-necked Grebe 5 June was a first for Churchill (S. Shadick) while a Pied-billed Grebe 30 June was an accidental there (BC). There were good indications of a Great Blue Heron colony at Ft. McMurray, n. of the published breeding range (JG).


An ad. Snowy Egret 19 June at Ninette was a Manitoba rarity (ph., DF, HL). A Least Bittern 30 June was another first for Churchill (BC). Five Brant 7 June and three on 12 June were accidentals there (BC). A Snow Goose at Whitebear, Sask., regularly associated with a Canada Goose in a small flock of Canada until all left about mid-June (S. Jordheim). A Garganey was photographed near Strathmore 12 June Alberta's third record (ML). Another accidental at Churchill was a Harlequin Duck 24 June (BC), while the first Ruddy Duck for the area was a 3 June (K. John).

HAWKS THROUGH COOTS Sharp-shinned Hawks were noticeably down in Saskatchewan's boreal forest (WH) while one wandered to Churchill 15 June (V. Chartier, D. Paulson). In Saskatchewan, Swainson's Hawks had more failures than usual; 35 of 71 nests in three areas produced 68 young. The late May snowstorm apparently caused six Golden Eagle nests to slump down clay cliffs. Prairie Falcons had average success and Merlins continued to reoccupy sites in the Kindersley-Kerrobert area deserted during the heavy dieldrin period of the 1960's (SH). A flock of $20\pm$ Swainson's Hawks in a field at Lyleton, Man., was unusual for 19 June (DF, HL). Marsh Hawks increased in the Raymore and Calgary areas (WH, SL, AW). Four young Peregrine Falcons were released in Saskatoon; two are known to have died and two have disappeared (BG). An adult sighted near Delisle, Sask., was unusual for 8 June (BG). A pair of Sandhill Cranes was found regularly s. of Water Valley, Alta., from late May through 5 June, $60\pm$ mi. s. of their published breeding range (L. Gibb, JP, LO, SJ). A white Am. Coot was seen with normally plumaged birds at Moose Jaw 15 July (S.&G. Wait, C. Letkeman).

SHOREBIRDS — On 14 July Chris Adam (ph.) discovered two ad. Piping

Plovers (plus Semipalmateds) with downy young on the s. shore of Lak Athabasca — a n. extension of it Saskatchewan published breedin range by more than 400 mi. He ha seen an adult in the same location 1 July 1981. Two ad. Mountain Plove were observed near Onefour, Alta., July (CH). An estimated 6,000 Rudd Turnstones were found near Churchill June (BC) while a Willet there 28 Jun (TS, DM) and 3 Purple Sandpipers 1 June were accidentals (BC). Whimbrel turned up at Birds Hill P.F near Winnipeg 1 July, an unusual dat (R. Koes). Rare shorebirds in th Calgary area included a Red Knot 2 July, a White-rumped Sandpiper 2 July, two W. Sandpipers 24 July, 1 Hudsonian Godwits 14 July and one c two Ruffs 24-31 July (JS, MS, RBa RBu, AS, B. Storm). Another Wester was seen near Cochrane, Alta., 3 Jun (SJ, LO). On 27 July a flock of 11 Marbled Godwits was found of Ingebright L., Sask., just w. of the Grea Sand Hills, as were 1,700 Wilson' Phalaropes (CH). What appeared to be a late non-breeding concentration of 3 Am. Avocets was noted 5 June nea Calgary (RBu). Pakowki L., Alta., had one Black-necked Stilt 15 May (G Thorn) and two 3 Aug. (ML). What is ap parently the second spring record for a Red Phalarope in Alberta was a full plumaged, studied for $10\pm$ minutes of a mudflat with other shorebirds, 3 June near Ft. McMurray (JG).

JAEGERS THROUGH HUMMING BIRDS — The first documented Sask atchewan record of a Long-tailed Jaeger, diagnosed as being in its second-year nuptial plumage, stayed a Last Mt. L. 8-17 July and was caugh (SL, WH, BK, ph., B. Luterbach); there are fewer than 10 records for the province. Alberta's third record for a Black-legged Kittiwake was an adult- o nearly adult-plumaged individual n.e. o Calgary 6 June (ph., JS, MS, W Wilson). Churchill's first California Gul



ong-tailed Jaeger

vas seen 8 and 11 June (F. Cooke) and s first Franklin's Gull 17 July (L&CW). accidentals there included a ther Black-headed Gull 15 July (DM, TS), a aughing Gull 9-25 July (BC, DM), Little Fulls periodically through the period vith a maximum of three 5 June (DM) nd an ad. Black-legged Kittiwake 18 uly (BC, m.ob.); Ross' Gulls peaked at 0 on 20 June (BC). Southern Manitoba ad its second Great Black-backed Jull, an immature, at Oak Hammock 2-June (T. ffrench, m.ob.). Stragglers ontinued at Churchill with a Black uillemot 17&18 July (BC, H. Currie). askatchewan's fifth Band-tailed ligeon was seen 25 and 28 June near laymore (WH). Great Horned Owls in askatchewan showed the effects of the eclining snowshoe hare population Iter peaking in 1981. About 30% of airs did not attempt nesting and 78 ests produceed 153 young, just under 0 young per successful nest (SH). urrowing Owls appeared to be up in anitoba and the Saskatoon area but own near Regina (H. Copland, BG, K). More than 50 breeding pairs settld in s. Manitoba where over 100 young ere banded (B. Ratcliff fide W. Koonz). W. C. Harris

Although nests of Pygmy and Boreal Owls are rarely found one of the former held two young near Priddis, Alta., 11 July (AW, KJ, GS); one of the latter had two young 25 July in Calgary (AS) and there was another nest at Birds Hill P.P., Man. (G. Smith). Late Com. Nighthawks were migrating 1 June (30 birds) at Good Spirit Lake and 2 June $(45\pm)$ at Regina (J.&W. Anaka, T. Riffel, BK). A very dark swift with a slightly forked tail, apparently a Black Swift, not previously reported in Saskatchewan, was studied for 5 minutes over Mortlach 4 July at heights from 30 to 80 ft. (P. Browne). A hummingbird with a completely orange body, the description of a Rufous Hummingbird, was seen 13 June n.w. of Southend, Sask., more than 500 mi. n.e. of its breeding range (K. Roney).

WOODPECKERS THROUGH BLUEBIRDS — A Lewis' Woodpecker strayed to Calgary 17 July (RBa). With about 15 Manitoba reports to date, a Scissor-tailed Flycatcher was a rarity at Oak Hammock 5 June (ph., DF). A Say's Phoebe 18 June was the first record for Churchill (K. Kaufman). Four species, recently common at Ft. McMurray, increased significantly over 1981: Tree and Barn Swallows more than doubled and Black-billed Magpies and Common Crows increased fourfold (JG). Tree Swallow nesting around Saskatoon was delayed until after the May storm, and was thus more uniform than usual; occupancy and success were down modestly with 91 successful and 21 unsuccessful nesting attempts. Nevertheless, the usually evident surplus of swallows seemed not to be present; it may be that half the swallow population perished during the storm (MH). The first summer record of a Black-capped Chickadee for Churchill was 11 June (BC). A breeding record for White-breasted Nuthatch was a rarity for Calgary (AS). The first recent breeding record for Sage Thrashers in Saskatchewan was a flightless young caught 8 July n.w. of Govenlock (ph., WH). Two Hermit Thrushes singing 17-23 July near Dundurn, Sask., were s. of their published breeding range (BG).



Sage Thrasher

W. C. Harris

WARBLERS THROUGH FINCHES — A Magnolia Warbler singing in Calgary 30 June was apparently out of its breeding range (JP) as were a 12 June Chestnut-sided, also singing (RBa), and a Cape May in the foothills w. of Calgary 18 July (AW, C.&M. Wershler). Connecticut Warblers an parently on territory were n. of the published breeding range at Ft. Md Murray in June (one bird; JG) and s. of near Priddis 11 July (2; AW, KJ, GS and near Bottrel, Alta., (at least 5; SJ Brewer's Blackbirds, another commo bird at Ft. McMurray, quadrupled thi summer (JG). A pair of Scarle Tanagers nesting at Dauphin, Man., wa n. of their previously known breeding range (B. Walley). In Calgary three sing ing Rose-breasted Grosbeaks 20 Juni hinted at breeding s. of their norma range (AS). An Indigo Bunting nea Roche Percee, Sask., 5 July indicated new breeding area (WH). Single Lazu Buntings were well out of range at Maid stone 20 June (WH) and near Beause jour, Man., 3 June (M. Siepman, G Grieef, N. Cleveland). There has been only one record of a Purple Finch a Churchill until one was seen 18 July (L&CW). Lone Sharp-tailed Sparrows singing 12 June near Irricana and Langdon, Alta., (JS, AW) and 18 July near Calgary (W. Hall) were beyond their normal breeding range. A Dark eyed Junco was seen in Moose Jaw 2⁻ June (M. Ritchie). A Field Sparrov showed up again this year near Beause jour, Man., through June and July (P Taylor). Clay-colored Sparrow numbers more than doubled at Ft. McMurray possibly because of increasing habita (JG). An undocumented Brewer's Sparrow was reported at Churchill 20 June, far from its closest range in s.w Saskatchewan (DM, J. Skykoff, TS).

INITIALED CONTRIBUTORS – (Provincial compilers in boldface; loca compilers in italics) – R. Barkley, *R Butot, B. Chartier*, **H. Copland**, D. Fast **B. Gollop**, J. Gulley, C. Harris, W Harris, M. Houston, S. Houston, *S Johnston*, K. Jones, **R. Koes**, B. Kreba S. Lamont, H. Lane, M. Lewis, D. McRae, L. Orman, J. Podlubny, G Simpson, T. Sinclair, A. Slater, *J. Steeves*, M. Steeves, L.&C. Weseloh, *A. Wiseley.*