CHECK-LIST OF SASKATCHE WAN MOTHS PART 7: SUMACH MOTHS (EUTELIINAE), SARROTHRIPIN MOTHS (SARROTHRIPINAE), NOLIN MOTHS (NOLINAE), MIDGETS (ACONTIINAE), PANTHEIN MOTHS (PANTHEINAE)

RONALD R. HOOPER, Saskatchewan Museum of Natural History, 2340 Albert Stree Regina, Saskatchewan. S4P 3V7

In the following text these abbreviations are used: s = south, n = north, w = west, e = east, CNC = the only Saskatchewan records of the species that we know of are in the Canadian National Collection in Ottawa. Unless otherwise indicated all the species are represented in the collection of the Saskatchewan Museum of Natural History, Regina.


EUTELIINAE - Sumach Moths

These are small moths 25 - 35 mm in wing span. Superficially they resemble Pyralid Moths (Pyralidae), but with broader bodies, and lacking elongated palpi. The hind wing often has a flattened lower edge instead of being even rounded. At rest the forewings a wrapped around the hind wings, wi abdomen elevated. Most species feed of the Sumach family (Anacardiaceae). O species feeds on Poison Ivy (*Rh radicans*).

Eyed Paectes - *Paectes oculatrix* (Gn.) Tantallon.

Expected species

Dark Marathyssa - *Marathyssa infici* (Wlk.) - e Canada, w to Manitoba ar British Columbia



Eyed Paectes

Lorie Melit, S.M.N.

SARROTHRIPINAE - Sarrothripin Moths

These are small moths 22 - 34 mm in wing span with long fore-wings which are broadened at the base giving a slightly rounded fore-edge as in Tortricid Moths (Tortricidae). The caterpillars feed on deciduous trees and shrubs.

Doubleday's Baileya - Baileya doubledayi (Gn.) - Tantallon.

Eyed Baileya - *B. ophthalmica* (Gn) - se Saskatchewan, w to Punnichy; n to Love.

rigid Owlet - *Nycteola frigidana* (Wlk.) central Saskatchewan,s to Whitewood area, n to Otter Rapids area.

NOLINAE - Nolin Moths

Small grey or white moths with a postnedian band of raised scales on the upper side of the fore-wings. The males have "feathered" antennae.

Confused Meganola - *Meganola minuscula* (Zell.) - Tantallon and Maryfield.

lurry-patched Nola - *Nola cilicoides* (Grt.) - s Saskatchewan,n to Wadena.

ACONTIINAE - Midgets

These are relatively small moths with a wing span from 16 to 30 mm. They have the head usually rounded in front with the elongated palpi upturned. The hindwings are usually unmarked. A number of the species have a contrasting black and white pattern on the fore-wings. When they are wrapped around the body at rest it provides protection by giving the insect the appearance of a bird dropping. Most of the caterpillars are semi-loopers because they lack some of their prolegs.

- Bog Lithacodia *Lithacodia bellicula* Hbn. - Earl Grey, Saskatoon, Carrot River, Aylsham and Canwood.
- Common Midget L. albidula (Gn.) n Saskatchewan, s to Northgate, Fort Qu'Appelle and Unity.
- Black-dotted Lithacodia L. synochitis (G. & R.) - Roche Percee and Rockglen.
- Pink-barred Lithacodia *L. carneola* (Gn.) - s Saskatchewan, n to Erwood and Aylsham.

nk-barred Lithacodia

Lorie Melit, S.M.N.H.

(4). December 1990



Yellow-cloaked Midget

Lorie Melit, S.M.N.F

- Pale-tipped Midget *Capis curvata* Grt. -Tantallon and Carrot River.
- Brown Leuconycta *Leuconycta lepidula* (Grt.) - e Saskatchewan, w to Buffalo Pound Provincial Park, n to Cumberland House.
- Yellow-cloaked Midget *Tarachidia semiflava* (Gn.) - s Saskatchewan, n to Prince Albert.
- Prairie Bird-dropping Moth T. binocula (Grt.) - s Saskatchewan, n to Wadena and Saskatoon.
- Olive-shaded Bird-dropping Moth T. tortricina (Zell.) - s Saskatchewan, n to Fort Qu'Appelle and Saskatoon.
- Arizona Bird-dropping Moth Conochares arizonae (Hy.Edw.) - Killdeer, Val Marie and Swift Current.

- Narrow-winged Midget *Therasea augus tipennis* (Grt.) - s Saskatchewan, n t Aylsham.
- Yellow-costa Midget *T. flavicosta* (Sm.) Rockglen, Val Marie and Tompkins.
- Sutrix Bird-dropping Moth Ponometi sutrix (Grt.) Saskatoon (CNC).

Expected Species:

- Pitcher-plant Moth Exyra rolandian Grt. - w to Aweme, Manitoba. The larv feeds on Pitcher-plant (Sarracenia pui purea).
- Fasciatella Midget *Fruva fasciatell* (Grt.) - n to Aweme, Manitoba.
- Common Spragueia *Spragueia leo* (Gn - nw to Aweme, Manitoba. The larve feeds on Bindweed.

PANTHEINAE - Panthein Moths

These are medium-sized moths. The males have "feathered antennae." The eyes are usually hairy as they are also in the Hadenin Moths (Hadeninae). The larvae have dense tufts of hair and feed on trees.

Panthea species - near *Panthea acronyctoides* (Wlk.) - central Saskatchewan, n to La Ronge and Meadow Lake Provincial Park, also in Cypress Hills.

Pale Panthea - *P. pallescens* McD. - Loon Lake.

Yellowhorn - *Colocasia flavicornis* (Sm.) - Duck Mountain Provincial Park, Hudson Bay and Greenwater Provincial Park.

lose-banded Yellowhorn - *C. propin-quilinea* (Grt.) - Bainbridge (n of Hudson Bay) and Fort Qu'Appelle.

The Laughter - *Charadra deridens* (Gn.) -Maryfield and Tantallon.

The Brother - *Raphia frater* Grt.- s Saskatchewan, n to La Ronge.

Expected Species:

Colorado Raphia - *R. coloradensis* Putnam-Cramer - n to Medicine Hat area of Alberta.

¹ BROWER, J.H. and A.E. BROWER 1970. Notes on the biology and distribution of moths associated with the pitcher plant in Maine. Proc. of Entomol. Soc. Ontario. Vol. 101:79-83.

² JONES, R.J.L. 1951. An annotated check list of the Macrolepidoptera of British Columbia. Occ. Paper Entomol. Soc. B.C., No. 1.

he Brother

Lorie Melit, S.M.N.H.

6(4). December 1990